

# CALF MILK REPLACER

RELIANCE CALF MILK REPLACER is an all milk powder blend providing excellent calf response to the natural food sources of milk fat, milk proteins and sugar.

## FEEDING RECOMMENDATIONS

Mixing ratio – 125g RELIANCE CALF MILK REPLACER to one litre of water

Dissolve RELIANCE CALF MILK REPLACER powder in hot (not boiling) water. Mix to a creamy, smooth consistency and add cool water to make up the required amount. Feed at body temperature (38-40°C).

Mix only when required.

Feed after a good intake of colostrum has been achieved (from four days after birth). Ensure that clean drinking water is available at all times.

Ensure that calves have access to hay and Reliance calf feed. Keep buckets, bottles and other equipment clean.

## FEEDING PROGRAM: ONCE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS PER DAY	LITRES / FEED	CONCENTRATION	DAILY INTAKE OF PELLETS/MUESLI (GRAMS)
1 - 4	2	2.5	Colostrum Only	-
5 - 14	2	2	125 grams / litre	50 - 100
15 - 28	1	2.5	250 grams / litre	100 - 350
29 - weaning	1	2.5	250 grams / litre	350 - 1500

## FEEDING PROGRAM: TWICE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS PER DAY	LITRES / FEED	CONCENTRATION	DAILY INTAKE OF PELLETS/MUESLI (GRAMS)
1 - 4	2	2.5	Colostrum Only	-
5 - 14	2	2	125 grams / litre	50 - 150
15 - 28	2	2.5	125 grams / litre	100 - 350
29 - weaning	2	2.5	125 grams / litre	350 - 1500

## INGREDIENTS SELECTED FROM

Milk Powder, Vitamin A, D3, E, K3, B1, B2, B3, B5, B6, B12, Ascorbic Acid, Potassium, Lasalocid (bovatec), Niacin, Pantothenic Acid, Biotin, Cobalt (organic), Copper (organic), Dextrose, Ferrous Iron, Iodine, Selenium (organic), Magnesium, Manganese (organic), Zinc (organic), Endox, Lecithin.



## TYPICAL ANALYSIS (As Fed Basis)

Crude Protein	25-26%
Crude Fat	19-20%
Lactose	44%
Minerals	6%

## COCCIDIOSTAT

RELIANCE CALF MILK REPLACER with BOVATEC®; registration number A009679

**FOR ANIMAL TREATMENT ONLY:**  
Contains Bovatec 20CC® at 500mg/kg of feed (100mg/kg of lasalocid)

**INDICATIONS:** Coccidiosis control in cattle, calves and goats.

**DIRECTIONS:** Feed as directed.

**Contraindications:** Care must be exercised when feeding concurrently with other antimicrobials.

Do not feed with other ionophores e.g monensin capsules, liquid or premix

Ensure recommended dose rates are not exceeded


Not to be used as a single dose treatment.

Do not feed to dogs, horses or other equids as ingestion by these species may be fatal.

### WITH-HOLDING PERIODS:

MEAT: Cattle and Goats – nil;

MILK: Cattle - Nil, Goats – 35 days.



**WARNING: NOT FOR HUMAN CONSUMPTION**



## You can rely on Reliance

# RELIANCE FEEDS



### NELSON TO NORTH OTAGO & WEST COAST

- Reliance Feed Mill Rolleston, 9 Link Drive
- Phone: 03 347 3670 or Fax: 03 347 3679
- Email: crtfeedrolleston@crt.co.nz

### SOUTH / CENTRAL OTAGO & SOUTHLAND

- Feed Mill Winton, 39 Great North Road
- Phone: 03 236 6014 or Fax: 03 236 7293
- Email: crtfeedwinton@crt.co.nz

OR CONTACT YOUR LOCAL CRT FARMLANDS FARMCENTRE OR TECHNICAL FIELD OFFICER



0800 278 583 [www.crtfarmlands.co.nz](http://www.crtfarmlands.co.nz)